

Lata Mangeshkar Medical Foundation's

Deenanath Mangeshkar Hospital & Research Center

Erandawane, Pune 411 004, Tel., 020 40151000 / 49153000

Email: info@dmhospital.org. Website: www.dmhospital.org



Joint Replacement and Reconstruction Discharge Summary

Patient Name: Mr. DARAKH DINESH

MRD#: 86258

SHANKARLAL

Sex: Male

Date Of Birth: 02/09/1962

Visit Code: IP0001

DOD: 16/09/2024

DOA: 12/09/2024

Room/Bed No.: SS8A - 3803

Consultant: Dr. WAKANKAR HEMANT Reference: DR DESHPANDE ATUL Diagnosis: Osteoarthritis of both knees.

Clinical course and Event:

C/O:Pain in both knees, difficulty in walking, climbing stairs.

K/C/O:NO COMORBIDITIES

O/E:Vitals - Normal, RS - Clear, CVS - NAD, P/A - Soft.

L/E:No Effusion, No PFT, MJLT + +, ROM R 10° - 120° L - 5° - 130°, Hips - Normal, No DNVC.

ASA Grade:

2

Pre op Hb: 14.4

Pre op PCV: 42.7

Pre op Sr Creat: 1.25

Pre op HbA1C: 6.3

Pre op EF: 64%

Post op Hb: 11

Post op PCV: 33

Post op Sr Creat: 0.9

Advice on Discharge:

1. T. PACIMOL 650 mg 1-1-1-1 x 7 days and then 1-1-1 x 7 days then SOS -60

2. T. PAN D 1-0-0 x 7 days - 7

3. T. ULTRACET SEMI 1-0-1 x 7 days and then 0-0-1 x 7 days then SOS

4.T. RESNER PLUS 0-0-1 x 1 month -3

5. T. XARELTO 10 mg 0-0-1 x 14 days -) *\

6. T. TRICIUM ACTIVE 1-0-0 x 2 months. — 60

7. T. ECOSPRIN 75 0-0-1 x 3 months(To be started only after course of T. XARELTO is over) Following medicines to be continued as per Physician's advice

TAB STAMLO 2.5 MG IF BP > 150/90 MM HG?

	KNEE ROM ON DISCHARGE	
RT	0	90
LT	Ü	9()

Removal on: 26/09/2024 مر

physiotherapy for 1 month after discharge: Physiotherap.
Next follow up at Decnanath Mangeshkar Hospital, Pune 3 MONTHS

With: Dr. Hemant Wakankar

OPD Time: Mon and Wed 3 pm

to 5 pm

By prior Appointment only. For appointment call 020-40151100 between 9 AM - 6 PM.

For emergency contact: Dept. ON Call no: 9850285871.

1165.

In case of emergency please come to DMH ER 1 (Emergency Room). It is open 24 hours a day. Phone no.: 020-4015-1027 / 1065.

OPERATIVE NOTE

Pre-operative Antibiotics:

Inj. SUPACEF 1.5gm IV

Operative Procedure: Bilateral Total Knee Replacement

Date of Surgery:13/9/24 Type of Anaesthesia:SA+LA Anaesthetist:Dr. SAWANT SHILPA

Tourniquet inflated at 280 mm Hg for Rt - 53 min, LT - 35 min.

Midline incision. Arthrotomy through medial para patellar approach.

Patella: No degeneration. Patellar osteophytes excised.

ACL- Attenuated. PCL- Normal. Menisci- Degenerated.

Routine medial and posteromedial release done.

Distal femoral cut taken using intramedullary jig at 6°. Tibial cut with posterior 7° slope using extramedullary jig. Femoral sizing done using anterior referencing and femoral cutting block fixed

Flexion / Extension gaps balanced and mediolateral stability checked. Posterior release done. in 3° of external rotation.

Anterior and posterior chamfer cuts and box cuts completed. Trial prosthesis inserted. Joint stability

patellofemoral tracking satisfactory. Lateral retinacular release not done. Femoral canal plugged with bone.

Trial prosthesis removed. Thorough joint wash given with normal saline pulsed lavage.

Tibial baseplate, Femoral prosthesis fixed with Palacos bone cement (Supacef 750 mg added).

Tourniquet released. Haemostasis achieved.

Tibial Poly inserted. Component stability and mediolateral stability reviewed.

Wound closed in layers without drain with Vicryl 1 and undyed Vicryl 2-0 and Staples.

The same procedure repeated on the other side.

Sterile dressing done.

DVT stockings given immediately post operatively.

Intraoperative PCV not given.

Inj. Tranexamic Acid (20 mg/kg) given at cementation and repeated at 3hrs and 6hrs.

LMWH started on the night of the surgery and continued till adequately mobilized.

Prosthesis Used - PERSONA, ZIMMER, USA

I EROOT III, EITHER					
Size	-	Right	Left		
Femur		6	6		
Tibia		D	D		
Poly		11 mm MC	10	mm	MC

Patella

Post operative Course.

Post operative pain controlled WITH LOCAL infiltration.

Inj. Supacef 750 mg IV X 2 doses. ALL physiotherapy goals achieved.

Foleys cathether: No ICU STAY: No

Blood Tranfusion: No

List of Emergencies:
Fever.swelling.bleeding.redness, calf swelling, tightness,pain. Please come to hospital for urgent

- 1. Not to wet the wound after surgery till sutures are removed.
- 2. Routine physiotherapy as per advice.
- 5. Do not start antibiotics on suspicion of infection without consulting the operating surgeon. Report 3. Start walking outdoor by 3 weeks.
- to Deenanath Hospital joint replacement department.
- 6. Review with Dr. Hemant Wakankar after 3 months with prior appointment.
- 7. Zuprinor Skin patch applied on 13/9/2024. Do not apply heat in area of Zuprinor skin patch application. Remove patch on 20/7/2024. Remove patch early in case of excessive nausea, vomiting or giddiness

Special needs

Physiotherapy:

For 1 month at home after surgery

Pain Relief:

Pain killers for 2 weeks, then as and when required

To be changed if soaked more than 50% in fresh blood

Date: 16/09/2024 Name of Dr: Prithviraj

Time: 10 am Designation: Registrar

Please bring this card for Follow up: Investigation attached separately: